

Feeding ecology of a nesting population of the Common Buzzard (*Buteo buteo*) in the Upper Nitra Region, Central Slovakia

Potravná ekológia hniezdnej populácie myšiaka hôrneho (*Buteo buteo*) na hornom Ponitri, stredné Slovensko

Karol ŠOTNÁR & Ján OBUCH

Šotnár K & Obuch J 2009: Feeding ecology of a nesting population of the Common Buzzard (*Buteo buteo*) in the Upper Nitra region, Central Slovakia. Slovak Rapt J 3: 13–20.

Abstract: During routine checks of the nests of the Common Buzzard in the Upper Nitra region (Central Slovakia) we collected food remains and recorded all prey given to the nestlings. We present results from the period 2006–2008. 606 food items were determined, comprising mainly mammals (67%). Birds were also frequent (17%), less so amphibians and reptiles (7%) and invertebrates (10%). During the rodent population peak in 2007, Buzzard pairs nesting in the valley preyed mainly on *Microtus arvalis*, whereas pairs living at the rim of the valley fed on *Myodes glareolus* and *Talpa europaea*, and higher in the Vtáčnik Mts birds fed on *Apodemus flavicollis*. After a massive decline in the rodent population in the following year 2008, the survival rate of the nestlings was very low and the proportion of invertebrates in their diet increased. Birds that were most frequently preyed upon included juvenile *Turdus philomelos* and *Garrulus glandarius*, prey identified from amphibian was mainly *Bufo bufo*, and from reptile prey was largely *Anguis fragilis*.

Abstrakt: Pri kontrolách hniezd myšiaka hôrneho v oblasti horného Ponitria (stredné Slovensko) sme zbierali zvyšky jeho potravy a zaznamenávali korisť, prinesenú mláďatám. Prezentujeme výsledky z rokov 2006–2008. Determinovali sme 606 kusov potravy, v ktorej dominujú cicavce (67 %). Početné sú aj vtáky (17 %), obojživelníky a plazy (7 %) a bezstavovce (10 %). Pri vysokej populačnej hladine hlodavcov v roku 2007 bol u párov, hniezdiacich v kotline, dominantnou korisťou druh *Microtus arvalis*, na okraji kotliny druhy *Myodes glareolus* a *Talpa europaea* a u párov, hniezdiacich vyššie v pohorí Vtáčnik druh *Apodemus flavicollis*. Po silnom poklese populačnej hladiny hlodavcov v nasledujúcom roku 2008 bola nízka úspešnosť hniezdenia myšiakov a v ich potrave stúpol podiel bezstavovcov. Z vtákov boli najčastejšie lovené mláďatá druhov *Turdus philomelos* a *Garrulus glandarius*, z obojživelníkov *Bufo bufo* a z plazov *Anguis fragilis*.

Key words: *Buteo buteo*, diet, Upper Nitra region, Central Slovakia

Karol Šotnár, Gavloviča 1/5, SK–971 01 Prievidza, Slovakia. E-mail: karol_sotnar@post.sk.

Ján Obuch, Comenius University, Botanical Garden, SK–038 15 Blatnica, Slovakia. E-mail: obuch@rec.uniba.sk.

Acknowledgments: We owe our gratitude to Martin Dúbrava and Branislav Beniska who physically examined the Buzzard nests, collected pellets and food remains of the Common Buzzard in the Upper Nitra region.

Introduction

In contrast to owls, bones are digested to a greater degree in the pellets of birds of prey. Therefore the investigation of their diet requires employing several different methods. During the nesting period, examination of pellets, determination of bone and feather remains in the nest can be used, as well as recording prey items presented to the nestlings. Prey remains in the nests of the *Buteo buteo* were evaluated by Šotnár (2003, 2007). In South Moravia this method was used for examining the food of the *B. buteo* by Voríšek et al. (1997) and Horák & Matušík (2000), a smaller sample from the Czech Republic and

Moravia was examined by Toufar (1958). In Germany, the same method was used to examine the food of *B. buteo* by Wuttke (1963) and Mebs (1964).

In the past when birds of prey were hunted legally and shot individuals were concentrated in taxidermist workshops, the stomach content of the birds could be examined. Sládek (1961, 1966) used this method in cooperation with the taxidermist workshop in Žiar nad Hronom. In the same manner, Sabadoš & Sládek (1968) studied the diet of the birds of prey shot in the pheasant reserve during an exceptionally harsh winter. Ševčík (1980) examined the food of the Buzzards shot in the pheasant reserves in the

Czech Republic. Extensive material collected by Rörig from Germany was studied by Uttendorfer (1939).

Comprehensive data from Central Europe is summarized in the papers of Glutz et al. (1989) and Cramp & Simmonds (1980). Our aim was to evaluate and compare the diet of the Common Buzzard population in three nesting seasons and three elevation zones of the Upper Nitra region.

Material and Methods

In 2006–2008 we collected pellets and recorded food remains in the nests of the Common Buzzard in the Upper Nitra region. Nests were examined in May–July. Our work builds upon the study by Šotnár & Topercer (2009). Collected material was classified into two categories:

Provisions – visual examination of food items. Determination of whole individuals, food remains, feather and fur directly in the nest.

Pellets – analysis of pellets and bone remains. All samples were processed in 5% NaOH solution to dissolve the feather and hair material. Bone fragments were used for the determination of vertebrate prey and elytrae and jaws were used to determine invertebrates. Mammals were determined according to cranial fragments, moles were also identified using humerus, Microtids using the lower molar teeth M_1 . Birds were determined according to 5 types of bone fragments: beaks, *humerus*, *metatarsus* and *tarsometatarsus*. Reptiles were determined according to jaws, skeletons and skin or scale remains. Food samples collected in 2007 and 2008 were classified into 3 zones:

A. Nesting pairs from the valley, coline zone 300–500 m a. s. l. (agroecosystems, field forests, riparian vegetation, 42 samples);

B. Nesting pairs from the piedmont area of Vtáčnik Mts, submontane zone 500–700 m a. s. l. (forest edge up to 1 km deep into the forest, 36 samples);

C. Nesting pairs from the mountain range Vtáčnik Mts., montane zone 700–1100 m a.s.l. (deep in the forests, approximately 2–5 km from the rim of the valley, 15 samples).

To evaluate the results in the Table 1–4 the method of marked differences from the mean (MDFM; Obuch 2001) was used. In the tables samples are sorted according to their similarity. Species in the upper part of the table are sorted so that those with marked positive deviations are arranged in blocks bordered by solid lines. Species with less marked deviations are sorted under the dashed line according to descending frequency. Those with frequency less than 5 are shown in Table 1. The lower part of the table shows sums for individual vertebrate and invertebrate classes. The diversity index H' is calculated according to the formula of Shannon & Weaver (1949).

Results

In 2006–2008 we studied food remains in the nests of *B. buteo* and determined 606 prey items. The most dominant part of the diet comprised mammals (Mammalia, 67%), less frequent were birds (Aves, 17%), amphibians and

Tab. 1. Food composition of *B. buteo* in the Upper Nitra Valley, inter-annual comparison

Tab. 1. Potrava *B. buteo* na hornej Nitre, porovnanie rokov

year (no.) / rok (č.) species / druhy	2006 (1)	2007 (2)	2008 (3)	Σ	%
<i>Apodemus flavicollis</i>	1- 1	1+ 35	1- 0	36	5.94
Coleoptera sp.	5	20	1+ 19	44	7.26
Hymenoptera sp.		1 1+	7	8	1.32
<i>Microtus arvalis</i>	38	110	1- 13	161	26.57
<i>Talpa europaea</i>	27	65	21	113	18.65
<i>Myodes glareolus</i>	6	37	9	52	8.58
<i>Turdus philomelos</i>	5	11	1	17	2.81
<i>Garrulus glandarius</i>	5	6	5	16	2.64
<i>Sorex araneus</i>		10	3	13	2.15
<i>Bufo bufo</i>	1	9	1	11	1.82
<i>Anguis fragilis</i>	3	2	3	8	1.32
<i>Coccothraustes coccothraustes</i>	4	2	2	8	1.32
<i>Sturnus vulgaris</i>	1	5	1	7	1.16
<i>Arvicola terrestris</i>	2	4	1	7	1.16
<i>Buteo buteo</i>		2	4	6	0.99
<i>Columba livia domestica</i>	1	2	2	5	0.83
<i>Terricola subterraneus</i>	2	3		5	0.83
Mammalia	78	275	1- 53	406	67.00
Aves	26	54	21	101	16.67
Amphibia, Reptilia	13	23	5	41	6.77
Evertebrata	8	1-	24 1+ 26	58	9.57
Σ	125	376	105	606	100.00
H' diversity index / index diverzity	2.49	2.58	2.67	2.74	

Other species / Ostatné druhy (year no. / rok č. – number / počet): *Erinaceus concolor* (1–1; 3–1), *Sorex minutus* (1–1), *Neomys fodiens* (2–1; 3–1), *Crocidura suaveolens* (2–1), *Lepus europaeus* (2–3; 3–1), *Sciurus vulgaris* (3–2), *Mus cf. musculus* (2–1), *Apodemus sylvaticus* (2–1), *Microtus agrestis* (2–1), *Mustela nivalis* (2–1), *Artiodactyla* sp. (2–2; 3–1), *Columba oenas* (3–1), *Columba palumbus* (1–1; 2–1; 3–1), *Columba* sp. (2–2), *Aegolius funereus* (2–1), *Strix aluco* (1–1), *Dryocopus martius* (2–1), *Dendrocopos medius* (2–1), *Dendrocopos* sp. (2–4; 3–1), *Anthus* sp. (2–1), *Locustella fluviatilis* (1–1), *Turdus* sp. (2–1), *Sitta europaea* (2–2), *Emberiza citrinella* (3–1), *Fringilla coelebs* (1–1; 3–1), *Pica pica* (2–1), *Passeriformes* sp. (1–6; 2–11; 3–1), *Bombina variegata* (1–1), *Rana temporaria* (1–1; 2–1), *Rana cf. esculenta* (2–1), *Amphibia* sp. (1–3), *Lacerta agilis* (2–1; 3–1), *Natrix natrix* (2–2), *Elaphe longissima* (1–1; 2–4), *Coronella austriaca* (1–1), *Colubridae* sp. (1–2; 2–3), *Orthoptera* sp. (1–3; 2–1), *Melolontha melolontha* (2–1), *Diptera* sp. (2–1)

reptiles (Amphibia and Reptilia, 7%; Fig. 1–7). Invertebrates (Evertebrata, 10%) are numerous, but comprise an insignificant part of the total biomass.

The most dominant prey species *Microtus arvalis* (26.6%) was less frequent in 2008, *Talpa europaea* (18.7%) is equally frequent in samples from 2006, 2007 and 2008, *Apodemus flavicollis* (5.9%) is more frequent in 2007. The lack of rodents in 2008 is partially compensated by increased proportion of invertebrates, including mainly insects of the orders Coleoptera and Hymenoptera (Table 1). The most frequent mammals include *Myodes glareolus* (8.6%) and *Sorex araneus* (2.2%). From birds *B. buteo* fed most frequently on juveniles of *Turdus philomelos* (2.8%) and *Garrulus glandarius* (2.6%). *Bufo bufo* (1.8%) was among the most frequent amphibian prey and *Anguis fragilis* (1.3%) was the most frequent reptile.

Tab. 2. Food composition of *B. buteo* in the Upper Nitra Valley, comparison between sampling methods

Tab. 2. Potrava *B. buteo* na hornej Nitre, porovnanie metód zberov

species / druhy sample type / druh zberu	a	b	Σ	%
<i>Buteo buteo</i>	1-	0	1+	6
				6 0.99
<i>Turdus philomelos</i>	1-	4	1+	13
				17 2.81
<i>Coccothraustes coccothraustes</i>	1-	0	1+	8
				8 1.32
<i>Sturnus vulgaris</i>	1-	1	1+	6
				7 1.16
<i>Garrulus glandarius</i>	2-	1	1+	15
				16 2.64
<i>Passeriformes</i> sp.	2-	0	2+	18
				18 2.97
<i>Apodemus flavicollis</i>		36	2-	0
				36 5.94
<i>Myodes glareolus</i>		52	2-	0
				52 8.58
<i>Microtus arvalis</i>		144	1-	17
				161 26.57
<i>Coleoptera</i> sp.	—	42	1-	2
				44 7.26
<i>Talpa europaea</i>	—	92	—	21
				113 18.65
<i>Sorex araneus</i>		13		
				13 2.15
<i>Bufo bufo</i>		11		
				11 1.82
<i>Anguis fragilis</i>		7		1
				8 1.32
<i>Hymenoptera</i> sp.		8		
				8 1.32
<i>Arvicola terrestris</i>		7		
				7 1.16
<i>Terricola subterraneus</i>		5		
				5 0.83
<i>Columba livia dom.</i>		5		
				5 0.83
<i>Elaphe longissima</i>		3		2
				5 0.83
Mammalia	363	1-	43	406 67.00
Aves	2-	17	2+	84 101 16.67
Amphibia, Reptilia		32		9 41 6.77
Evertebrata		55	1-	3 58 9.57
Σ	467		139	606 100.00
H' diversity index / index diverzity	2.35		2.81	2.74
a – pellets / vývržky; b – nest supply / zásoba na hniezde				

The pellets revealed three times more prey items than food provisions in the nests. Therefore the comparison of species composition using the MDFM method (Table 2) showed marked differences only in the less frequent material collected in the nests. The pellets comprised mainly mammals (77.7%), including *M. arvalis*, *M. glareolus* and *A. flavicollis*. Coleopterans were the most frequent invertebrate prey. Nest provisions included mainly birds (Aves, 60.4%), namely *G. glandarius*, *T. philomelos*, *Sturnus vulgaris* and *Coccothraustes coccothraustes*.

The evaluation of *B. buteo* diet composition in relation to habitat was strongly influenced by the gradation of rodents in 2007 and the subsequent decline combined with unfavourable weather conditions in 2008. Great abundance of food in 2007 was reflected in the higher nesting success. Therefore in 2007 we collected a three times larger sample of food than in 2008. In 2007 the most frequent prey item of *B. buteo* in the valley included *M. arvalis* (51.4%), in the piedmont of Vtáčnik, *M. glareolus* and *T. europaea* were more frequent in the montane zone, *A. flavicollis* was dominant. In 2008, Aves and Evertabrata were more frequent in the valley and *T. europaea* and Evertebrata in the piedmont area. The montane zone yielded only a small sample of 22 food items. Notable prey items included *Aegolius funereus*, *Strix aluco* and *Locustella fluviatilis*.

Discussion

Our data on the food composition of *B. buteo* from the Upper Nitra valley, which were collected through the analysis of pellets and food provisions in the nests, corresponds with the results of other authors using similar methods in South Bohemia and Germany. Since the works of Voříšek et al. (1997) and Horák & Matušík (2000) only include prey determined to higher taxonomic levels, we decided to make comparison in Table 4 only with our own data from the determination of pellets at Pálava. All four studies on the diet of *B. buteo*, used in the comparison, show a lower proportion of the dominant species *M. arvalis* and higher proportion of the subdominant *T. europaea*. Birds can be seen in higher proportion, especially *G. glandarius* and *T. philomelos* species. In contrast to other studies, our results show higher proportion of *S. araneus*, *A. flavicollis*, *C. coccothraustes* and *Elaphe longissima*. In our results we also included findings of dead *B. buteo* nestlings, killed mainly as a result of Cainism during the shortage of food in 2008. Studies using similar methods found other species more abundant such as *Apodemus sylvaticus*, *Arvicola terrestris*, *Lacerta agilis*, *A. fragilis* (Mebs 1964) *Lepus europaeus*, *Alauda arvensis* (Mebs 1964, Wuttky 1963), *Cricetus cricetus*,

K. Šotnár



Fig. 1. The most frequently preyed bird – *Garrulus glandarius* (Veľká Čausa, 29 May 2008, 1 pull. 13-day and 1 egg).

Obr. 1. Najčastejšie lovený druh vtákov bol *Garrulus glandarius* (Veľká Čausa, 29. máj 2008, 1 pull. 13-dňové a 1 vajce).

K. Šotnár



Fig. 3. Food supply – *T. europaea*, *A. terrestris*, *L. fluviatilis*, *T. merula* (Lehotá pod Vtáčnikom, 6 June 2006, pull. 21 days old).

Obr. 3. Zásoba potravy – *T. europaea*, *A. terrestris*, *L. fluviatilis*, *T. merula* (Lehotá pod Vtáčnikom, 6. jún 2006, mláďa 21-dňové).

K. Šotnár



Fig. 5. Main food component of the Common Buzzard – *Microtus arvalis* (Nováky, 6 June 2007, 2 pull. ca 28 days old).

Obr. 5. Hlavná zložka potravy myšiakov – *Microtus arvalis* (Nováky, 6. jún 2007, 2 pull. asi 28-dňové).

K. Šotnár




Fig. 2. Prey item in the nest – *Sciurus vulgaris* without head (Fančová, 27 May 2008, 2 pull. 5–7 days old).

Obr. 2. Potrava na hniezde – *Sciurus vulgaris* bez hlavy (Fančová, 27. máj 2008, 2 pull. 5–7-dňové).

K. Šotnár



Fig. 4. Prey item – *Coronella austriaca* (Cígel, Vtáčnik, 24 May 2004, 2 pull. ca 18–20 days old and 1 egg).

Obr. 4. Potrava – *Coronella austriaca* (Cígel, Vtáčnik, 24. máj 2004, 2 pull. asi 18–20 dňové a 1 vajce).

K. Šotnár



Fig. 6. Three nestlings in the nest, food supply – 6 *M. arvalis* (Kamenec p. Vtáčnikom, 24 May 2008).

Obr. 6. Tri mláďatá na hniezde, potrava na hniezde – 6 *M. arvalis* (Kamenec p. Vtáčnikom, 24. máj 2008).

Tab. 3. Food composition of *B. buteo* in the Upper Nitra Valley, evaluation of 2007 and 2008 samples according to habitat zones
Tab. 3. Potrava *B. buteo* na hornej Nitre, vyhodnotenie zberov z rokov 2007 a 2008 podľa zón

year / rok	2007						2008			Σ	%		
	A		B		C		A	B	C				
zone / zóna II species / druhy	1+	73	1-	34	1-	3	10	1-	2	1-	1	123	25.47
<i>Microtus arvalis</i>	1+	73	1-	34	1-	3	10	1-	2	1-	1	123	25.47
<i>Myodes glareolus</i>	2-	2	1+	26		9			3		6	46	9.52
<i>Talpa europaea</i>	1-	13	1+	44		8	1-	1	1+	18		86	17.81
<i>Apodemus flavicollis</i>	1-	5		15	1+	15						35	7.25
Coleoptera sp.	10		1-	6		4	1+	8		7	4	39	8.07
Hymenoptera sp.	—	1	—	—	—	—	—	2	1+	5	—	8	1.66
<i>Sorex araneus</i>	4	—	3	—	3	—			1	—	2	13	2.69
<i>Turdus philomelos</i>	2		7		2				1			12	2.48
<i>Garrulus glandarius</i>	3		3				3		1		1	11	2.28
<i>Bufo bufo</i>	2		5		2				1			10	2.07
<i>Buteo buteo</i>			2				4					6	1.24
<i>Sturnus vulgaris</i>	3		2				1					6	1.24
<i>Arvicola terrestris</i>	3		1				1					5	1.04
<i>Anguis fragilis</i>	2					1		1		1		5	1.04
Mammalia	106	129		40	1-	14		25		14		328	67.91
Aves	17	31		7	1+	15		4		3		77	15.94
Amphibia, Reptilia	7	11		5		1		3		1		28	5.80
Evertebrata	12	1-	8		4	1+	10	1+	12		4	50	10.35
Σ	142		179		56		40		44		22	483	100.00
H' diversity index / index diverzity	2.11		2.50		2.36		2.44		2.00		2.22		2.73

Zone: A – coline zone 300–500 m a. s. l., B – submontane zone 500–700 m a. s. l., C – montane zone 700–1100 m a. s. l.

Zóna: A – kolínna 300–500 m n. m., B – submontánna 500–700 m n. m., C – montánná 700–1100 m n. m.

Emberiza citrinella, *Columba livia domestica* (Wuttky 1963) and Coleoptera (Voříšek et al. 1997; Table 4).

The papers focusing on the examination of stomach contents more frequently studied winter samples, dominated mainly by *M. arvalis*. Stomach contents of Common Buzzards shot in the pheasant reserves more often included *Phasianus colchicus* and *Gallus gallus domesticus* species (Ševčík 1980). Invertebrates are usually better preserved in stomach samples than vertebrates, therefore Coleoptera, Orthoptera (Sládek 1961), as well as Lepidopteran larvae (caterpillars) were often found in the stomach contents (Ševčík 1980, Uttendorfer 1939).

According to our study from the Upper Nitra, small game (hare, pheasant, grey partridge) comprises less than 1% of the Common Buzzard's diet. Several authors (Sládek 1966, Horák & Matušík 2000) suggested that small game were not originally preyed upon by *B. buteo* and that it only fed on their carcasses.



Fig. 7. *Astacus* sp. – unusual prey of Common Buzzard. (Lehota pod Vtáčnikom, 30 May 2009).

Obr. 7. *Astacus* sp. – nezvyčajná potrava na hniezde myšiaka hôrneho. (Lehota pod Vtáčnikom, 30. máj 2009).

Tab. 4. Evaluation of data on *B. buteo* diet in the Central Europe
Tab. 4. Vyhodnotenie údajov o potrave *B. buteo* v strednej Európe

species / druhy	1	2	3	4	5	6	7	Σ	%
<i>Sorex araneus</i>	1+	13		3	4	7	2-	0	27
<i>Buteo buteo</i>	1+	6			1			7	0.11
<i>Coccothraustes coccothr.</i>	1+	8		1	2		1-	0	11
<i>Elaphe longissima</i>	1+	5						5	0.08
Hymenoptera sp.	1+	8						8	0.13
<i>Apodemus flavicollis</i>	2+	36	7	3	11	1-	0	57	0.93
<i>Talpa europaea</i>	2+	113	1+ 31	1+ 25	2-	14	1-	115	5.49
<i>Turdus philomelos</i>	2+	17	2+ 17	1	1-	0	2-	0	35
<i>Garrulus glandarius</i>	2+	16	1+ 6	2	1-	0	2-	0	24
<i>Bufo bufo</i>	1+	11	1+ 7	1	5	2	2-	0	26
<i>Sturnus vulgaris</i>	1+	7	1+ 7				1-	0	14
<i>Myodes glareolus</i>	2+	52	11	5	1+ 11	1-	12	4	95
<i>Apodemus sylvaticus</i>	1	1+ 7	4	3	10	1	2-	0	26
<i>Arvicola terrestris</i>	7	1+ 7		2	3			10	0.47
<i>Turdus merula</i>		1+ 6	4		3		2	1-	15
<i>Lacerta agilis</i>	2	1+ 15		3	1-	4	1-	0	30
<i>Anguis fragilis</i>	8	3+ 39		2	2-	1	1-	0	76
<i>Lepus europaeus</i>	1-	4	1+ 21	1+ 11	6	16	6	33	1.58
<i>Alauda arvensis</i>		1+ 8	1+ 7				1-	0	15
<i>Cricetus cricetus</i>	2-	0	1- 0	3+ 56		2-	3	1	57
<i>Emberiza citrinella</i>	1			1+ 6		2	1	1-	10
<i>Columba livia dom.</i>	5			1+ 9		1	5	2-	0
Coleoptera sp.	45	3-	0	19	3+ 79	1+ 137	1+ 69	5-	1
<i>Gryllotalpa gryllota</i>	1-	0	1- 0		2+ 58	1- 0	2-	7	65
Orthoptera sp.	2-	4	2- 0	2- 0	2+ 221	2- 3	3-	12	240
<i>Gallus gallus dom.</i>				2		3	1+ 7	4	16
<i>Phasianus colchicus</i>					1	1-	1	1- 11	40
Lepidoptera sp.	3-	0	3- 0	2- 0	2- 0	4- 0	2+ 27		0.65
<i>Microtus arvalis</i>	161	1-	156	2- 61	1- 48	587	323	1+ 172	259
<i>Terricola subterraneus</i>	—	5	— 1	— 2	— 2	— 2	— 1	— 0	52.66
<i>Perdix perdix</i>	—	—	1	— 2	— 5	— 1	18	27	0.44
<i>Lacerta vivipara</i>			5		3			10	0.29
<i>Oryctolagus cuniculus</i>				3			1	12	0.26
<i>Mustela nivalis</i>	1		1		1	1	1	10	0.24
<i>Capreolus capreolus</i>			2	4				4	10
<i>Rattus norvegicus</i>			3			1	2	3	0.15
<i>Sciurus vulgaris</i>	2		1		1	1		4	0.15
<i>Natrix natrix</i>	2							6	0.13
Pisces sp.						1	1	6	0.13
<i>Rana temporaria</i>	2		4					6	0.10
<i>Corvus corone</i>			3			1	2	6	0.10
<i>Fringilla coelebs</i>	2			3	1			6	0.10
<i>Microtus agrestis</i>	1			1		3	1	6	0.10
<i>Neomys fodiens</i>	2					1	3	6	0.10
<i>Mus cf. musculus</i>	1					4		5	0.08
<i>Carduelis carduelis</i>				1	4			5	0.08
Mammalia	406	266	215	1- 105	1- 735	376	2248	4351	70.89
Aves	1+ 101	1+ 48	2+ 72	17	79	1+ 87	2- 39	443	7.22
Amphibia, Reptilia, Pisces	41	1+ 70	2- 2	1- 6	2- 26	3- 5	1+ 264	414	6.74
Vertebrata	1- 58	4- 0	1- 19	1+ 79	1+ 418	1+ 160	1- 196	930	15.15
Σ	606	384	308	207	1258	628	2747	6138	100.00
H' diversity index / index diverzity	2.74	2.38	2.65	2.06	2.07	1.93	1.36	2.31	

1 – own data / vlastné údaje, 2 - Mebs 1964, Germany / Nemecko, 3 - Wuttky 1963, Germany / Nemecko, 4 – Voríšek et al. 1997, south Moravia / južná Morava, 5 – Sládeček 1961, Slovakia / Sovaňsko, 6 – Ševčík 1980, Bohemia / Čechy, 7 – Uttendorfer 1939, Germany / Nemecko

Tab. 5. Comparison of *B.buteo* food composition with diet of *A. gentilis* and *A. pomarina* in Central Slovakia
Tab. 5. Porovnanie potravy *Buteo buteo* s potravou *Accipiter gentilis* a *Aquila pomarina* na strednom Slovensku

species / druhy	Apom		Bbut		Agen		Σ	%
<i>Microtus arvalis</i>	1+	394	1-	161	4-	4	559	37.62
<i>Microtus agrestis</i>	1+	40	2-	1	1-	0	41	2.76
<i>Arvicola terrestris</i>	1+	57	2-	7	2-	0	64	4.31
<i>Mustela nivalis</i>	1+	11		1			12	0.81
<i>Rana temporaria</i>	1+	13		2			15	1.01
<i>Talpa europaea</i>	2-	16	1+	113	3-	0	129	8.68
<i>Sorex araneus</i>	1-	0	1+	13			13	0.87
<i>Apodemus flavicollis</i>	1-	9	1+	36		5	50	3.36
<i>Myodes glareolus</i>	3-	0	1+	52	1-	1	53	3.57
<i>Bufo bufo</i>	1-	0	1+	11			11	0.74
Coleoptera sp.	2-	3	1+	45	2-	0	48	3.23
Hymenoptera sp.			1+	8			8	0.54
<i>Columba livia dom.</i>	3-	4	3-	5	2+	94	103	6.93
<i>Garrulus glandarius</i>	2-	1		16	2+	31	48	3.23
<i>Buteo buteo</i>	1-	0		6	1+	8	14	0.94
<i>Phasianus colchicus</i>		1			1+	6	7	0.47
<i>Streptopelia decaocto</i>	1-	0	1-	0	1+	13	13	0.87
<i>Dendrocopos major</i>	1-	0	1-	0	1+	12	12	0.81
<i>Turdus merula</i>		1	1-	0	1+	11	12	0.81
<i>Turdus philomelos</i>	2-	1		17	1+	20	38	2.56
<i>Sciurus vulgaris</i>	1-	0		2	1+	9	11	0.74
<i>Coccothraustes coccothraustes</i>	1-	0		8		6	14	0.94
<i>Lepus europaeus</i>		5		4		2	11	0.74
<i>Anguis fragilis</i>		3		8			11	0.74
<i>Sturnus vulgaris</i>		1		7		2	10	0.67
<i>Columba palumbus</i>		2		3		3	8	0.54
<i>Perdix perdix</i>		2				5	7	0.47
<i>Strix aluco</i>				1		5	6	0.40
<i>Columba oenas</i>				1		4	5	0.34
<i>Terricola subterraneus</i>				5			5	0.34
<i>Elaphe longissima</i>				5			5	0.34
Mammalia	1+	541		406	3-	22	969	65.21
Aves	2-	34	1-	101	2+	260	395	26.58
Amphibia, Reptilia		19	1+	41	2-	0	60	4.04
Evertebrata	2-	4	1+	58	2-	0	62	4.17
Σ		598		606		282	1486	100.00
H' diversity index / index diverzity		1.55		2.74		2.72	2.81	

Apom – *Aquila pomarina* (Dravecký et al. 2008, central Slovakia / stredné Slovensko), **Bbut** – *Buteo buteo* (own data / vlastné údaje),
Agen – *Accipiter gentilis* (Šotnár 2000)

A study of the diet of two similar sized birds of prey; *Accipiter gentilis* (Šotnár 2000) and *Aquila pomarina* (Dravecký et al. 2008) carried out in Upper Nitra region were studied in the Upper Nitra region: compare their diets (Tab. 5). In *A. pomarina* greater proportion of the diet comprises *M. arvalis* and several other non-sylvatic vole species: *M. agrestis* and *A. terrestris*. The most frequently

hunted frog species include *R. temporaria*, whereas *B. buteo* hunts *B. bufo* more frequently. *A. gentilis* is markedly specialized to hunting birds (92% of the diet), few mammals killed include mainly *Sciurus vulgaris*. Block comparison of species comprising the majority of the diet shown in Tab. 5 reveals that the food niches of these 3 larger bird of prey species have a minimal overlap.

References

- Cramp S & Simmons KEL (eds) 1980: The Birds of the Western Palearctic, Vol. II., Hawks to Busards. Oxford Univ Press, Oxford, London & New York. 696.
- Dravecký M, Danko Š, Obuch J, Kicko J, Maderič B, Karaska D, Vrána J, Šreibr O, Šotnár K, Vrlík P & Bohačík L 2008: Diet of the Lesser Spotted Eagle (*Aquila pomarina*) in Slovakia. Slovak Rapt J 2: 1–18.
- Glutz von Blotzheim UN, Bauer KM & Bezzel E 1989: Handbuch der Vögel Mitteleuropas. Bd. 4 Falconiformes [2., durchgesehene Auflage]. Aula Verlag, Wiesbaden, 943.
- Horák P & Matušík H 2000: Ještě k potravě káně lesní [Notes on the diet of Common Buzzard]. Crex 16: 33–37. [In Czech with English summary]
- Mebs T 1964: Zur Biologie und Populationsdynamik des Mäusebussards (*Buteo buteo*). J Ornithol 105(3): 247–306.
- Obuch J 2001: Using marked differences from the mean (MDFM) method for evaluation of contingency tables. Buteo 12: 37–46.
- Sabadoš K & Sládek J 1968: Príspevok k poznaniu potravy dravcov a sov v bažantnici počas mimoriadne tuhej zimy 1962–1963 [Beitrag zur Kenntnis der Nahrung der Greifvögel und der Eulen in einer Fasaneerie während des aussergewöhnlich strengen Winters 1962–1963]. Biológia 23 (5): 363–369. [In Slovak with German summary]
- Shannon CE & Weaver W 1949: The mathematical theory of communication. Univ Illinois Press, Urbana, 125.
- Sládek J 1961: Príspevok k poznaniu potravnej ekológie myšiaka lesného *Buteo buteo* (L.) [Beitrag zur Kenntnis der Nahrungsökologie des Mäusebussards *Buteo buteo* L.]. Zool listy, 10(1): 331–343. [In Slovak with German summary]
- Sládek J 1966: Poznanie zákonitostí potravnej ekológie ako základ hodnotenia hospodárskeho významu myšiaka lesného (*Buteo buteo*) [Erkenntnis der Gesetzmäßigkeiten der Ernährungsökologie als Grundlage der wirtschaftlichen Bedeutung des Mäusebussards (*Buteo buteo*)]. Českoslov Ochr Prírody 12: 93–103. [In Slovak with German and Russian summaries]
- Ševčík J 1980: Potrava káně lesní (*Buteo buteo*), káně rousné (*Buteo lagopus*) a jestřába lesního (*Accipiter gentilis*) v oblastech s intenzivním chovem bažantů [Die Ernährung des Mäusebussards (*Buteo buteo*), Rauhfussbussards (*Buteo lagopus*) und Habichts (*Accipiter gentilis*) in der Gegend mit einer Fasanenzucht]. Sylvia 20: 35–43. [In Czech with German summary]
- Šotnár K. 2000: Príspevok k hniezdnej biológii a potravnej ekológiu jastraba veľkého (*Accipiter gentilis*) na hornom Ponitri [A contribution to the breeding biology and feeding ecology of the Goshawk (*Accipiter gentilis*) in the Horné Ponitrie Region]. Buteo 11: 43–50. [In Slovak with English summary]
- Šotnár K 2003: Poznatky o hniezdnej populácii myšiaka hôrneho (*Buteo buteo*) [Knowledge on breeding population of Common Buzzard (*Buteo buteo*)]. Vtácie správy 11(1): 17. [In Slovak]
- Šotnár K 2007: Dva roky spod hniezd myšiakov hôrnych [Two years below the nests of Common Buzzard]. Dravce a sovy 3(1): 18–19. [In Slovak with English summary]
- Šotnár K & Topercer J 2009: Estimating density, population size and dynamics of Common Buzzard (*Buteo buteo*) in a West Carpathian region by a new method. Slovak Rapt J 3: 1–12.
- Toufar J 1958: Příspěvek k poznání potravy mladých kání lesních (*Buteo buteo*) podle zbytků kořisti na hnizdech [Zur Kenntnis der Nahrung nestjunger Mäusebussarde (*Buteo buteo*)]. Sylvia 15: 67–76. [In Czech with German and Russian summaries]
- Uttendorfer O 1939: Die Ernährung der deutschen Raubvögel und Eulen und ihre Bedeutung in der heimischen Natur. Verlag J Neumann, Neudamm, 412.
- Voříšek P, Krištín A, Obuch J & Votýpka J 1997: Potrava káně lesní v České republice a její význam pro myslivost [Diet of Common Buzzard in the Czech Republic and its importance for gamekeeping]. Buteo 9: 57–68. [In Czech with English summary]
- Wuttke K 1963: Beutetier-Funde in Greifvögelhorsten des Hakel. Beitr Vogelkd 93(1/2, 3): 140–171.